

Mathieu series approach to the Hilbert inequality

Tibor K. Pogány

Faculty of Maritime Studies, University of Rijeka, 51000 Rijeka, Croatia
Institute of Applied Mathematics, Óbuda university, 1034 Budapest, Hungary
poganj@pfri.hr

Using the concept of the Mathieu $(\mathbf{a}, \boldsymbol{\lambda})$ -series a set of generalizations are derived for the discrete Hilbert inequality, when the kernel function is non-homogeneous.

References

- [1] B. Drašćić Ban, T. K. Pogány, Discrete Hilbert type inequality with non-homogeneous kernel, *Appl. Anal. Discr. Math* **3** (2009), No. 1, 88-96.
- [2] B. Drašćić Ban, J. Pečarić, T. Pogány, On a discrete Hilbert type inequality with non-homogeneous kernel, *Sarajevo J. Math.* **6** (2010), No. 1, 23-34.
- [3] B. Drašćić Ban, J. Pečarić, I. Perić, T. Pogány, Discrete multiple Hilbert type inequality with non-homogeneous kernel, *J. Korean Math. Soc.* **47** (2010), No. 3, 537-546.
- [4] D. Jankov, T. K. Pogány, R. K. Saxena, Extended general Hurwitz–Lerch Zeta function as Mathieu $(\mathbf{a}, \boldsymbol{\lambda})$ -series, *Appl. Math. Letters* **24** (2011), No. 8, 1473-1476.
- [5] T. K. Pogány, Integral representation of Mathieu $(\mathbf{a}, \boldsymbol{\lambda})$ -series, *Integral Transforms Spec. Functions* **16** (2005), 685–689.
- [6] T. K. Pogány, Integral representation of a series which includes the Mathieu \mathbf{a} - series, *J. Math. Anal. Appl.* **296** (2004), No.1, 309-313.
- [7] T. K. Pogány, Hilbert’s double series theorem extended to the case of non-homogeneous kernels, *J. Math. Anal. Appl.* **342**(2008), No. 2, 1485–1489.
- [8] T. K. Pogány, New class of inequalities associated with the Hilbert’s double series theorem, *Appl. Math. E-Notes* **10** (2010), 47-51.
- [9] T. K. Pogány, H. M. Srivastava, Ž. Tomovski, Some families of Mathieu \mathbf{a} -series and alternating Mathieu \mathbf{a} -series, *Appl. Math. Comput.* **173**(2006), No. 1, 69-108.