

Univalence criteria related with the generalised Sălăgean and Ruscheweyh operator

Páll-Szabó Ágnes Orsolya

Babeş-Bolyai University, Cluj-Napoca, Romania

pallszaboagnes@math.ubbcluj.ro

In this paper we obtain new sufficient conditions for univalence using the method of Loewner chains using the generalised Sălăgean and Ruscheweyh operator. In particular, we obtain some known univalence conditions due to Lewandowski, Becker, Kanas and Lecko.

References

- [1] J. Becker, *Löwnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen*, J. Reine Angew. Math. 255 (1972), 23-43.
- [2] P. L. Duren, H. S. Shapiro, A. L. Shields, *Singular measures and domains not for Smirnov type*, Duke Math. J. 33 (1966), 242-254.
- [3] S. Kanas, A. Lecko, *Univalence criteria connected with arithmetic and geometric means*, II, Proceedings of the Second Int. Workshop of Transform Methods and Special Functions, Varna '96, Bulgar. Acad. Sci. (Sofia) (1996), 201-209.
- [4] Z. Lewandowski, *On a univalence criterion*, Bull. Acad. Polon. Sci. Ser. Sci. Math., 29 (1981), 123-126.
- [5] A. E. Nistor-Şerban, *Univalence criteria related with Sălăgean and Ruscheweyh operators*, Bulletin of the Transilvania University of Brasov Vol 9(58), No. 2 - 2016 Series III: Mathematics, Informatics, Physics, 41-48.
- [6] Pommerenke, Ch., *Über die Subordination analytischer Funktionen*, J. Reine Angew Math., 218(1965), 159-173.
- [7] G. S. Sălăgean, *Subclasses of univalent functions*, Lecture Notes in Math. (Springer Verlag), 1013(1983), 362-372.
- [8] St. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc., 49(1975), 109-115.