

On Laplacian eigenvalues of graphs

Shariefuddin Pirzada

Department of Mathematics, University of Kashmir, India

`pirzadasd@kashmiruniversity.ac.in`

Let $G(V, E)$ be a simple graph of order n , size m and having the vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. The adjacency matrix $A = (a_{ij})$ of G is a $(0, 1)$ -square matrix of order n whose (i, j) -entry is equal to 1 if v_i is adjacent to v_j and equal to 0, otherwise. Let $D(G) = \text{diag}(d_1, d_2, \dots, d_n)$ be the diagonal matrix associated to G , where $d_i = \deg(v_i)$, for all $i = 1, 2, \dots, n$. The matrix $L(G) = D(G) - A(G)$ is called the Laplacian matrix and its spectrum is called the Laplacian spectrum (L -spectrum) of the graph G . We discuss some of the recent developments of Laplacian eigenvalues and Laplacian energy of graphs.

References

- [1] H. Bai, The Grone-Merris conjecture, *Trans. Amer. Math. Soc.* **363** (2011) 4463–4474.
- [2] A. E. Brouwer and W. H. Haemers, Spectra of graphs. Available from: <http://homepages.cwi.nl/aeb/math/ipm.pdf>.
- [3] Hilal A. Ganie, S. Pirzada, Rezwana and X. Li, Upper bounds for the sum of Laplacian eigenvalues of a graph and Brouwer's conjecture, *Discrete Math. Algorithms Applications*, 11, 2 (2019) 195008 (15 pages).
- [4] Hilal A. Ganie, A. M. Alghamdi and S. Pirzada, On the sum of the Laplacian eigenvalues of a graph and Brouwer's conjecture, *Linear Algebra Appl.* **501** (2016) 376-389.
- [5] S. Pirzada and Hilal A. Ganie, On the Laplacian eigenvalues of a graph and Laplacian energy, *Linear Algebra Appl.* **486** (2015) 454–468.